









## Monitoring Logging in Central Africa

## NASA- LCLUC 2002 Workshop






Dr. Nadine Laporte, Tiffany Lin, Didier Devers Department of Geography

University of Maryland, College Park, MD 20742 http://luci.umd.edu.umd/lcluc









Selective logging is the most extensive land use in the region, with millions of hectares of forested land under concession (i.e. allocated for logging). In order to better manage these forests for timber extraction and biodiversity conservation, we are using satellite imagery (30m Landsat and 1-4m IKONOS) and field surveys to monitor and analyze logging activities at several study sites in the region. Here we focus on the Tri-National Park area on the border between Cameroon, Central African Republic, and the Republic of Congo.

Our Integrated Forest Monitoring System (INFORMS) for logging includes: (A) maps of the distribution of logging activities in the region, (B) development of new integrated monitoring and , wildlife conservation, and carbon modeling activities, (D) quantification of deforestation rates around logging towns, and using satellite imagery. All of these results are conveyed to a newly created Central Africa Global Observations of Forest Cover



comparison to that in the Amazonian forests, with an average extraction rate of  $\pm 1$  tree/ha, or roughly 10 m<sup>3</sup> of wood/ha. With the exception of Gabon, most of the trees harvested are species of African mahogany (Entandrophragma spp.). Until recently, okoume (Aucoumea klaineana) had been the main species harvested in Gabon. It is now replaced by secondary species Ceiba pentadra, which is likely an indication of the impoverishment of the Gabonese forest. Together with adequate forest management policies and planning, accurate mapping of forest types & distribution and monitoring of land cover changes are essential to ensure the sustainability of timber resources.



Laporte N., (2001), Forest monitoring in Central Africa, Briefing sheets #13, Biodiversity Support Program, March 2001, Washington DC, 6p.

Data, International Society for Optical Engineering (SPIE), Toulouse, 17-21 September 2001, Toulouse, France, 6p.

Remote Sensing, 19(18): 3537-3550.

Rainforest Ecology and Conservation, Yale University Press, 119-138p.

Research funded by the

NASA/LCLUC & the


**USAID CARPE program** 

Laporte N., (2001), Tropical deforestation in Central Africa, Briefing sheets #6, Biodiversity Support Program, March 2001, Washington DC, 4p.

Laporte N., Goetz S.J., Justice C.O., Heinicke M., (1998), A new land cover map of central Africa derived from multi-resolution, multi-temporal AVHRR Data, International Journal


Wilkie D., Laporte N., (2001), Forest area and deforestation in Central Africa: Current knowledge and future directions, in W. Weber, A. Vedder, S. Morland, L. White (Eds), Africal

## **INFORMS Project** LOGGING MONITORING APPROACH Remote Sensing Analysis & Data Set Integration Forest Information Users & Collaborators National forest services Technical backstopping of OSFAC network Logging companies Remote sensing data acquisition and distribution Conservation programs Development of new validation methods (video/inventories) Research community • Remote sensing analysis & ecological field data integration Bringing expertise for field surveys: • Remote sensing/GIS training (Lope workshop 2000) Forest and fauna inventories • Primary study sites (Tri-National Park area - C.A.R, Rep. of Congo, and Cameroon; Okapi Reserve- D.R.C) Validation of forest maps Forest monitoring methods Forest type maps Biomass distribution Logging intensity estimates Results/ Outputs:



Bare soil and Roads (BS)

Classified Landsat TM 1999



**ESTIMATION OF LOGGING INTENSITY** 

timating forest logging intensity, CIB - Forest unit Mokobo 1998